

Special characters on star graphs and representations of $\ast$ -algebras

Vasyl Ostrovskiy*

Institute of Mathematics, National Acad. Sci. of Ukraine
E-mail: vo@imath.kiev.ua

Introduction

In a series of recent papers (see [4, 2, 13, 15] and references therein), families of self-adjoint operators $A_1, \dots, A_n$ were studied such that $A_1 + \dots + A_n = \lambda I$ and the spectrum of each operator A_l , $l = 1, \dots, n$, lies in fixed finite set M_l . Such families can be naturally regarded as $\ast$ -representation of certain algebra related to a star-shaped graph Γ and a positive function (character) on the set of its vertexes.

While the cases where Γ is a Dynkin graph or extended Dynkin graph were studied in details (see [2, 7, 6, 9] and others), in the case of general graph only partial results are known [4, 8, 12].

In the case of extended Dynkin graphs, well-known are special characters, which possess extra invariance properties. In this paper, we introduce such special characters for general star-shaped graphs and study their properties (Sec. 3). The formulas based on the positive solutions of certain equation, which is related to the graph (Sec. 2). Following [12] we decompose the special character into odd and even parts and study their evolution under reflections (Sec. 4). Notice that similar formulas for the evolution of the even and odd parts have been obtained independently in [11]. The obtained formulas are used in Sec. 5 to prove that for any graph containing an extended Dynkin graph as a proper subgraph there exists a character such that the corresponding algebra has infinite-dimensional irreducible $\ast$ -representation (the converse statement that for the case of extended Dynkin graph all irreducible representations are finite-dimensional, was proved in the previous paper [10]) Notice that this fact was formulated as a hypothesis in [12].

*This work was supported by the DFG, grant no. 436UKR 113/71/0-1 and State Foundation for Fundamental Research of Ukraine, grant no. 01.07/071

1 Preliminaries

Let Γ be a star-shaped graph, i.e., simply-laced non-oriented graph consisting of n branches, such that l -th branch has $k_l + 1$ vertexes, $l = 1, \dots, n$, and all branches are connected at a single root vertex (which is extreme point for all branches). Any function on the set of vertexes, is defined by a *character* on the graph,

$$\chi = (\alpha_1^{(1)}, \dots, \alpha_{k_1}^{(1)}; \dots; \alpha_1^{(n)}, \dots, \alpha_{k_n}^{(n)}; \lambda), \quad (1)$$

where $\alpha_1^{(l)}, \dots, \alpha_{k_l}^{(l)}$ are the values on the veriest of l -th branch starting from non-root extreme point, and λ is the value on the root vertex.

Given a graph Γ and a character χ on Γ consider the $*$ -algebra $\mathcal{A}_{\Gamma, \chi}$ over $\mathbb{C}$ generated by self-adjoint elements a_l , $l = 1, \dots, n$, which satisfy relations

$$p_l(a_l) = 0, \quad \sum_{l=1}^n a_l = \lambda e,$$

where $p_l(x) = x(x - a_1^{(l)}) \dots (x - a_{k_l}^{(l)})$, $l = 1, \dots, n$. Then $*$ -representations of this algebra are n -tuples $A_1, \dots, A_n$ of self-adjoint operators in a Hilbert space, such that $A_1 + \dots + A_n = \lambda I$ and the spectrum of each A_l is contained in $\{a_1^{(l)}, \dots, a_{k_l}^{(l)}\}$.

The essential tool to study $*$ -representations of $\mathcal{A}_{\Gamma, \chi}$ are reflection (Coxeter) functors introduced in [5] (for non-involutive case, see [1]). Namely there exist two functors, $S: \text{Rep } \mathcal{A}_{\Gamma, \chi} \rightarrow \text{Rep } \mathcal{A}_{\Gamma, \chi'}$ and $T: \text{Rep } \mathcal{A}_{\Gamma, \chi} \rightarrow \text{Rep } \mathcal{A}_{\Gamma, \chi''}$, where

$$\begin{aligned} \chi' &= (\alpha_{k_l}^{(1)} - \alpha_{k_l-1}^{(1)}, \dots, \alpha_{k_l}^{(1)} - \alpha_0^{(1)}; \dots; \alpha_{k_l}^{(n)} - \alpha_{k_l-1}^{(n)}, \dots, \alpha_{k_l}^{(n)} - \alpha_0^{(n)}; \lambda'), \\ \lambda' &= \alpha_{k_1}^{(1)} + \dots + \alpha_{k_n}^{(n)} - \lambda, \\ \chi'' &= (\lambda - \alpha_{k_1}^{(1)}, \dots, \lambda - \alpha_1^{(1)}; \dots; \lambda - \alpha_{k_n}^{(n)}, \dots, \lambda - \alpha_1^{(n)}; \lambda) \end{aligned}$$

(we put $\alpha_0^{(l)} = 0$, $l = 1, \dots, n$.) The action of these functors on $*$ -representations gives rise to the action on the set of characters, $S: \chi \mapsto \chi'$, $T: \chi \mapsto \chi''$. This action is extensively used to study the structure and properties of representations of algebras introduced above.

2 Equation related to a graph

Let n and k_l , $l = 1, \dots, n$ be given natural numbers, and let Γ be the corresponding star-shaped graph. In what follows, we will use solutions of the following equation

$$n - 1 - t = \sum_{l=1}^n \frac{1}{1 + t + \dots + t^{k_l}}, \quad t \geq 0. \quad (2)$$

Theorem 1. (See [10]). *The equation (2) have no solutions on $[0, \infty)$ if and only if the corresponding graph Γ is one of the Dynkin graphs A_d , $d \geq 1$, D_d , $d \geq 4$, E_6 , E_7 , or E_8 . The equation (2) has a unique solution on $[0, \infty)$ if and only if the corresponding graph Γ is one of the extended Dynkin graphs $\tilde{D}_4$, $\tilde{E}_6$, $\tilde{E}_7$, or $\tilde{E}_8$, this solution is $t = 1$. In all other cases (i.e., where Γ is neither a Dynkin graph nor an extended Dynkin graph), the equation (2) has on $[0, \infty)$ two solutions $0 < t_1 < 1 < t_2 < n - 1$.*

We show that $t_1 t_2 = 1$.

Proposition 1. *Let t be a solutions of (2). Then t^{-1} is also a solution of (2).*

Proof. Indeed,

$$\begin{aligned} t^{-1} + \sum_{l=1}^n \frac{1}{1 + t^{-1} + \dots + t^{-k_l}} &= t^{-1} \left(1 + \sum_{l=1}^n \frac{t^{k_l+1}}{1 + t + \dots + t^{k_l}} \right) \\ &= t^{-1} \left(1 + \sum_{l=1}^n \frac{t^{k_l+1}}{1 + t + \dots + t^{k_l}} + n - 1 - t - \sum_{l=1}^n \frac{1}{1 + t + \dots + t^{k_l}} \right) \\ &= t^{-1} \left(n - t + \sum_{l=1}^n \frac{t^{k_l+1} - 1}{1 + t + \dots + t^{k_l}} \right) = n - 1. \quad \square \end{aligned}$$

Hypothesis 1. *Let r be the largest eigenvalue of a star-shaped graph Γ . Then $t + t^{-1} + 2 = r^2$, where t is a positive solution of (2).*

3 Special characters

For graphs which are not Dynkin graphs, we introduce special character as follows:

$$\begin{aligned} \chi_\Gamma &= (\alpha_1^{(1)}, \dots, \alpha_{k_1}^{(1)}; \dots; \alpha_1^{(n)}, \dots, \alpha_{k_1}^{(n)}; 1), \\ \alpha_j^{(l)} &= \frac{1 + t + \dots + t^{j-1}}{1 + t + \dots + t^{k_l}}, \end{aligned} \quad (3)$$

where t is a solution of (2). For extended Dynkin graph such character is unique (since (2) has a unique solution $t = 1$) and has the form

$$\begin{aligned} \chi_{\tilde{D}_4} &= \frac{1}{2}(1; 1; 1; 1; 2); \\ \chi_{\tilde{E}_6} &= \frac{1}{3}(1, 2; 1, 2; 1, 2; 3); \\ \chi_{\tilde{E}_7} &= \frac{1}{4}(1, 2, 3; 1, 2, 3; 2; 4); \\ \chi_{\tilde{E}_8} &= \frac{1}{6}(1, 2, 3, 4, 5; 2, 4; 3; 6). \end{aligned}$$

For any graph containing extended Dynkin graph there are two special characters corresponding to two positive solutions of (2).

Proposition 2. $(TS)\chi_\Gamma = t^{-1}\chi_\Gamma$.

Proof. Indeed, the equality

$$\sum_{l=1}^n \frac{t^{k_l}}{1 + \dots + t^{k_l}} = n - 1 - \frac{1}{t}$$

implies that

$$\sum_{l=1}^n \frac{1 + \dots + t^{k_l-1}}{1 + \dots + t^{k_l}} = 1 + \frac{1}{t}, \quad (4)$$

and by a direct calculations we get $(TS)\chi_\Gamma = \frac{1}{t}\chi_\Gamma$. $\square$

Given a graph Γ consider the following quadratic form on the set of characters χ :

$$\gamma_\Gamma(\chi) = \sum_{l=1}^n \sum_{j=1}^{k_l} (\alpha_j^{(l)})^2 + \lambda^2 - \sum_{\substack{l=1 \\ k_l > 1}}^n \sum_{j=1}^{k_l-1} x_j^{(l)} x_{j+1}^{(l)} - \lambda \sum_{l=1}^n x_{k_l}^{(l)} \quad (5)$$

Proposition 3. For special character, $\gamma_\Gamma(\chi_\Gamma) = 0$.

Proof. We have, using (4)

$$\begin{aligned} \gamma_\Gamma(\chi_\Gamma, 1) &= \sum_{l=1}^n \sum_{j=1}^{k_l} \frac{(1 + \dots + t^{j-1})^2}{(1 + \dots + t^{k_l})^2} + 1 \\ &\quad - \sum_{\substack{l=1 \\ k_l > 1}}^n \sum_{j=1}^{k_l-1} \frac{(1 + \dots + t^{j-1})(1 + \dots + t^j)}{(1 + \dots + t^{k_l})^2} - \sum_{l=1}^n \frac{1 + \dots + t^{k_l-1}}{1 + \dots + t^{k_l}} \\ &= \sum_{l=1}^n \frac{1}{(1 - t^{k_l+1})^2} \left(\sum_{j=1}^{k_l} (1 - t^j)^2 - \sum_{j=1}^{k_l-1} (1 - t^j)(1 - t^{j+1}) \right) - \frac{1}{t}. \end{aligned}$$

Here we assume $t \neq 1$, since for $t = 1$ (Γ being an extended Dynkin graph) the statement is well known.

For $k_l > 1$ we have

$$\frac{1}{(1 - t^{k_l+1})^2} \left(\sum_{j=1}^{k_l} (1 - t^j)^2 - \sum_{j=1}^{k_l-1} (1 - t^j)(1 - t^{j+1}) \right) = \frac{1 - t^{k_l}}{(1 - t^{k_l+1})(1 + t)}.$$

For $k_l = 1$ the same holds assuming the second summand is zero. Then by (4) we have

$$\gamma_\Gamma(\chi_\Gamma) = \frac{1}{t+1} \sum_{l=1}^n \frac{1 - t^{k_l}}{1 - t^{k_l+1}} - \frac{1}{t} = 0. \quad \square$$

4 Evolution of odd and even components

In this section, similarly to [12] we decompose the special character into odd and even parts (in [12] these parts are called odd and even standard characters) and study their evolution under the action of the Coxeter functors. We obtain for arbitrary star-shaped graphs the formulas obtained in [12] for special classes of graphs.

As shown in [2], there is a correspondence between non-degenerate $*$ -representations of algebras $\mathcal{A}_{\Gamma, \chi}$ and non-degenerate locally scalar representations of Γ . Namely, $*$ -representations of the algebra $\mathcal{A}_{\Gamma, \chi}$ correspond to locally scalar representations of the graph Γ with character

$$u_{\Gamma} = (x_1^{(1)}, \dots, x_{k_1}^{(1)}; \dots; x_1^{(n)}, \dots, x_{k_n}^{(n)}; 1) \quad (6)$$

where

$$\begin{aligned} x_{k_l}^{(l)} &= \alpha_{k_l}^{(l)}, & x_{k_l-1}^{(l)} &= \alpha_{k_l}^{(l)} - \alpha_1^{(l)}, \\ x_{k_l-2}^{(l)} &= \alpha_{k_l-1}^{(l)} - \alpha_1^{(l)}, & x_{k_l-3}^{(l)} &= \alpha_{k_l-1}^{(l)} - \alpha_2^{(l)}, \\ x_{k_l-4}^{(l)} &= \alpha_{k_l-2}^{(l)} - \alpha_2^{(l)}, & x_{k_l-5}^{(l)} &= \alpha_{k_l-2}^{(l)} - \alpha_3^{(l)}, \end{aligned} \quad (7)$$

and so on; $l = 1, \dots, n$.

Applying these formulas for the special character χ_{Γ} we have the following special character of locally scalar representation:

$$\begin{aligned} x_{k_l}^{(l)} &= \frac{1 + t + \dots + t^{k_l-1}}{1 + t + \dots + t^{k_l}}, & x_{k_l-1}^{(l)} &= \frac{t(1 + t + \dots + t^{k_l-2})}{1 + t + \dots + t^{k_l}}, \\ x_{k_l-2}^{(l)} &= \frac{t(1 + t + \dots + t^{k_l-3})}{1 + t + \dots + t^{k_l}}, & x_{k_l-3}^{(l)} &= \frac{t^2(1 + t + \dots + t^{k_l-4})}{1 + t + \dots + t^{k_l}}, \end{aligned} \quad (8)$$

and so on.

Following [3] we decompose the set of vertexes Γ_v into odd and even parts, $\Gamma_v = \overset{\bullet}{\Gamma}_v \cup \overset{\circ}{\Gamma}_v$. However, unlike in [3], we always assume that the root vertex is odd. According to the decomposition of Γ_v we decompose u_{Γ} into odd and even parts, $u_{\Gamma} = \overset{\bullet}{u}_{\Gamma} + \overset{\circ}{u}_{\Gamma}$, where

$$\begin{aligned} \overset{\bullet}{u}_{\Gamma} &= (\dots, 0, x_{k_1-3}^{(1)}, 0, x_{k_1-1}^{(1)}, 0; \dots; \dots, 0, x_{k_n-3}^{(n)}, 0, x_{k_n-1}^{(n)}, 0; 1), \\ \overset{\circ}{u}_{\Gamma} &= (\dots, 0, x_{k_1-2}^{(1)}, 0, x_{k_1}^{(1)}; \dots; \dots, 0, x_{k_n-2}^{(n)}, 0, x_{k_n}^{(n)}; 0). \end{aligned}$$

Denote by σ_g the reflection at a point $g \in \Gamma_v$. Let $\overset{\bullet}{c}$ and $\overset{\circ}{c}$ be compositions of reflections at all odd and even points respectively (the order is irrelevant since all odd reflections commute and all even reflections commute). For detailed definitions and properties of these mappings, see [3].

Proposition 4. *Let the character of locally scalar representation be defined by (6), (8). For odd vertexes $\sigma_g(x_g) = t^{-1}x_g$. For even vertexes $\sigma_g(x_g) = tx_g$.*

Proof. Straightforward calculations. $\square$

Proposition 5. *The following formulas hold for all $j = 1, 2, \dots$*

$$\begin{aligned} (\overset{\bullet}{c}\overset{\circ}{c})^j \overset{\bullet}{u}_\Gamma &= \frac{1}{t^j(1-t)} ((1-t^{2j+1}) \overset{\bullet}{u}_\Gamma + t(1-t^{2j}) \overset{\circ}{u}_\Gamma), \\ \overset{\circ}{c}(\overset{\bullet}{c}\overset{\circ}{c})^j \overset{\bullet}{u}_\Gamma &= \frac{1}{t^j(1-t)} ((1-t^{2j+1}) \overset{\bullet}{u}_\Gamma + (1-t^{2j+2}) \overset{\circ}{u}_\Gamma), \\ (\overset{\circ}{c}\overset{\circ}{c})^j \overset{\circ}{u}_\Gamma &= \frac{1}{t^j(1-t)} ((1-t^{2j}) \overset{\bullet}{u}_\Gamma + (1-t^{2j+1}) \overset{\circ}{u}_\Gamma), \\ \overset{\bullet}{c}(\overset{\circ}{c}\overset{\circ}{c})^j \overset{\circ}{u}_\Gamma &= \frac{1}{t^{j+1}(1-t)} ((1-t^{2j+2}) \overset{\bullet}{u}_\Gamma + t(1-t^{2j+1}) \overset{\circ}{u}_\Gamma). \end{aligned}$$

Proof. The following fact follows from the Proposition 4 above.

Lemma 1. *Let u_Γ be defined by (6), (8). Then $\overset{\bullet}{c}\overset{\bullet}{u}_\Gamma = -\overset{\bullet}{u}_\Gamma$, $\overset{\circ}{c}\overset{\circ}{u}_\Gamma = -\overset{\circ}{u}_\Gamma$, $\overset{\circ}{c}\overset{\bullet}{u}_\Gamma = \overset{\bullet}{u}_\Gamma + (1+t)\overset{\circ}{u}_\Gamma$, $\overset{\bullet}{c}\overset{\circ}{u}_\Gamma = \overset{\circ}{u}_\Gamma + (1+t^{-1})\overset{\bullet}{u}_\Gamma$,*

Now the statement follows by induction from Lemma 1. $\square$

Remark 1. *Since $\overset{\bullet}{c}\overset{\bullet}{u}_\Gamma = -\overset{\bullet}{u}_\Gamma$ and $\overset{\circ}{c}\overset{\circ}{u}_\Gamma = -\overset{\circ}{u}_\Gamma$, Proposition 5 obviously gives the formulas for $(\overset{\circ}{c}\overset{\circ}{c})^j \overset{\bullet}{u}_\Gamma$, $\overset{\bullet}{c}(\overset{\circ}{c}\overset{\circ}{c})^j \overset{\bullet}{u}_\Gamma$, $(\overset{\circ}{c}\overset{\circ}{c})^j \overset{\circ}{u}_\Gamma$, $\overset{\circ}{c}(\overset{\circ}{c}\overset{\circ}{c})^j \overset{\circ}{u}_\Gamma$ as well.*

Corollary 1. *For a character of the form $v = \alpha \overset{\bullet}{u}_\Gamma + \beta \overset{\circ}{u}_\Gamma$ ($\alpha \neq 0$) introduce a “normalized” character $\tilde{v} = \overset{\bullet}{u}_\Gamma + \alpha^{-1} \beta \overset{\circ}{u}_\Gamma$. Then we have formulas similar to [12]:*

$$\begin{aligned} \widetilde{(\overset{\circ}{c}\overset{\circ}{c})^j \overset{\bullet}{u}_\Gamma} &= \overset{\bullet}{u}_\Gamma + \frac{t}{1+t} \rho_{\lambda-2}(2j) \overset{\circ}{u}_\Gamma, \\ \overset{\circ}{c} \widetilde{(\overset{\bullet}{c}\overset{\circ}{c})^j \overset{\bullet}{u}_\Gamma} &= \overset{\bullet}{u}_\Gamma + (1+t)(\rho_{\lambda-2}(2j+1))^{-1} \overset{\circ}{u}_\Gamma \\ &= \overset{\bullet}{u}_\Gamma + \frac{t}{1+t} (\lambda+2 - \rho_{\lambda-2}(2j)) \overset{\circ}{u}_\Gamma, \\ \widetilde{(\overset{\circ}{c}\overset{\bullet}{c})^j \overset{\circ}{u}_\Gamma} &= \overset{\bullet}{u}_\Gamma + (1+t)(\rho_{\lambda-2}(2j))^{-1} \overset{\circ}{u}_\Gamma \\ &= \overset{\bullet}{u}_\Gamma + \frac{t}{1+t} (\lambda+2 - \rho_{\lambda-2}(2j-1)) \overset{\circ}{u}_\Gamma, \\ \overset{\bullet}{c} \widetilde{(\overset{\circ}{c}\overset{\circ}{c})^j \overset{\circ}{u}_\Gamma} &= \overset{\bullet}{u}_\Gamma + \frac{t}{1+t} \rho_{\lambda-2}(2j+1) \overset{\circ}{u}_\Gamma. \end{aligned}$$

Here $\lambda = t + t^{-1}$ and following [12] we denote

$$\rho_{\lambda-2}(n) = 1 + \frac{t - t^n}{1 - t^{n+1}}.$$

We use the formula

$$(\rho_{\lambda-2}(n+1))^{-1} = \frac{t}{(t+1)^2} (\lambda+2 - \rho_{\lambda-2}(n)),$$

which is verified directly. Notice that in [12], the condition $t > 1$ was assumed, while we admit as t any of the two positive solutions of (2).

Proposition 6. *Let u'_Γ be the special character, constructed by the same formulas (6), (8) as u_Γ but with t^{-1} instead of t . Then for $t > 1$*

$$\begin{aligned}\lim_{j \rightarrow \infty} \widetilde{(\overset{\bullet}{c} \overset{\circ}{c})^j \overset{\bullet}{u}_\Gamma} &= \lim_{j \rightarrow \infty} \overset{\bullet}{c} \widetilde{(\overset{\circ}{c} \overset{\bullet}{c})^j \overset{\circ}{u}_\Gamma} = u_\Gamma, \\ \lim_{j \rightarrow \infty} \overset{\circ}{c} \widetilde{(\overset{\bullet}{c} \overset{\circ}{c})^j \overset{\bullet}{u}_\Gamma} &= \lim_{j \rightarrow \infty} \widetilde{(\overset{\circ}{c} \overset{\bullet}{c})^j \overset{\circ}{u}_\Gamma} = u'_\Gamma,\end{aligned}$$

and for $t < 1$

$$\begin{aligned}\lim_{j \rightarrow \infty} \widetilde{(\overset{\circ}{c} \overset{\circ}{c})^j \overset{\bullet}{u}_\Gamma} &= \lim_{j \rightarrow \infty} \overset{\bullet}{c} \widetilde{(\overset{\circ}{c} \overset{\circ}{c})^j \overset{\circ}{u}_\Gamma} = u'_\Gamma, \\ \lim_{j \rightarrow \infty} \overset{\circ}{c} \widetilde{(\overset{\circ}{c} \overset{\circ}{c})^j \overset{\bullet}{u}_\Gamma} &= \lim_{j \rightarrow \infty} \widetilde{(\overset{\circ}{c} \overset{\circ}{c})^j \overset{\circ}{u}_\Gamma} = u_\Gamma.\end{aligned}$$

Proof. Let $t > 1$. It was shown in [12] that $\lim_{j \rightarrow \infty} \rho_{\lambda-2}(j) = 1 + t^{-1}$ which implies the first formula.

The same formula yields

$$\lim_{j \rightarrow \infty} \overset{\circ}{c} \widetilde{(\overset{\bullet}{c} \overset{\circ}{c})^j \overset{\bullet}{u}_\Gamma} = \overset{\bullet}{u}_\Gamma + t \overset{\circ}{u}_\Gamma$$

Let $\overset{\bullet}{u}'_\Gamma$ and $\overset{\circ}{u}'_\Gamma$ be odd and even components of u'_Γ . The nonzero components of $\overset{\bullet}{u}_\Gamma$ have the form

$$x_{k_l-2j+1}^{(l)} = \frac{t^j(1+t+\dots+t^{k_l-2j})}{1+t+\dots+t^{k_l}}, \quad j = 0, 1, \dots$$

which is invariant under replacement $t \mapsto t^{-1}$. Therefore, $\overset{\bullet}{u}'_\Gamma = \overset{\bullet}{u}_\Gamma$.

The nonzero components of $\overset{\circ}{u}_\Gamma$ have the form

$$x_{k_l-2j}^{(l)} = \frac{t^j(1+t+\dots+t^{k_l-2j-1})}{1+t+\dots+t^{k_l}}, \quad j = 0, 1, \dots$$

which implies $\overset{\circ}{u}'_\Gamma = t \overset{\circ}{u}_\Gamma$.

For $t < 1$ we have $\lim_{j \rightarrow \infty} \rho_{\lambda-2}(j) = 1 + t$. The rest of the proof is the same. $\square$

5 Representations

Proposition 7. *Let Γ be a star-shaped graph. If Γ contains an extended Dynkin graph as a proper subgraph, then there exists a character χ on Γ such that $\mathcal{A}_{\Gamma, \chi}$ has infinite-dimensional irreducible $*$ -representation.*

Otherwise, all irreducible $$ -representations of $\mathcal{A}_{\Gamma, \chi}$ are finite-dimensional regardless of the choice of χ .*

Proof. It was shown in previous paper [10] that for Dynkin graphs and extended Dynkin graphs all irreducible $*$ -representations of $\mathcal{A}_{\Gamma, \chi}$ are finite-dimensional. Therefore, we need to prove the first part of the proposition only.

Let $\Gamma' \supset \Gamma$. Then any irreducible representation of $\mathcal{A}_{\Gamma, \chi}$ can trivially be extended to irreducible representation of $\mathcal{A}_{\Gamma', \chi'}$, where χ' coincides with χ on vertexes of Γ and is arbitrary outside Γ . Therefore, it remains to prove the statement for the following graphs: $\Gamma_{(1,1,1,1,1)}$, $\Gamma_{(1,1,1,2)}$, $\Gamma_{(2,2,3)}$, $\Gamma_{(1,3,4)}$, $\Gamma_{(1,2,6)}$, since any star-shaped graph containing an extended Dynkin graph as a proper subgraph, contains (or coincides with) one of the five listed above. Here we denote by $\Gamma_{(k_1, \dots, k_n)}$ the graph with n branches of lengths $k_1, \dots, k_n$.

After some calculations one can see that the components of the characters

$$w_p = \overset{\bullet}{u}_{\Gamma} + \frac{1 - t^p}{1 - t^{p-1}} \overset{\circ}{u}_{\Gamma},$$

are increasing along branches, $(w_p)_{j+1}^{(l)} > (w_p)_j^{(l)}$, for $p \geq k = \max k_l$. Let χ_p be the corresponding character on the graph constructed by the procedure inverse to (7). Then the components of χ_p are positive and increasing along branches, $(\chi_p)_{j+1}^{(l)} > (\chi_p)_j^{(l)}$, $j = 1, \dots, k_l - 1$, $l = 1, \dots, n$. This property of characters is essential for the constructions related to the corresponding algebra $\mathcal{A}_{\Gamma, \chi_p}$. Moreover, for $p = k$, we have $(\chi_k)_k^{(l)} = 1$ ($k_l = k$) which allows to construct for this algebra one-dimensional representation.

Since the action of T and S is induced by the action of $\overset{\circ}{c}$ and $\overset{\bullet}{c}$, applying powers of (TS) , to this representation, we get irreducible representations of the whole family of algebras $\mathcal{A}_{\Gamma, \chi_j}$, $j = p + 2r$, $r = 0, 1, \dots$.

Applying Proposition 6 and results of [14] we conclude that the algebra $\mathcal{A}_{\Gamma, \chi_r}$ has at least one $*$ -representation.

If $\mathcal{A}_{\Gamma, \chi_r}$ has a finite-dimensional irreducible representation, then the trace equality implies that 1 belongs to the rational span of components of χ_r . To complete the proof, we show that this is not true for the five graphs listed above. This is known for $\Gamma_{(1,1,1,1,1)}$ [4] and for $\Gamma_{(1,1,1,2)}$ [8]

Indeed, after simple transformations we find that t is a root of the corresponding polynomial $p_{\Gamma}(\cdot)$:

$$\begin{aligned} p_{\Gamma_{(1,1,1,1,1)}}(x) &= x^2 - 3x + 1, \\ p_{\Gamma_{(1,1,1,2)}}(x) &= x^4 - x^3 - x^2 - x + 1, \\ p_{\Gamma_{(2,2,3)}}(x) &= x^6 - x^4 - x^3 - x^2 + 1, \\ p_{\Gamma_{(1,3,4)}}(x) &= x^8 - x^5 - x^4 - x^3 + 1, \\ p_{\Gamma_{(1,2,6)}}(x) &= x^{10} + x^9 - x^7 - x^6 - x^5 - x^4 - x^3 + x + 1. \end{aligned}$$

It is a routine exercise to check that each of these polynomials is irreducible over $\mathbb{N}$ and therefore over $\mathbb{Q}$. Therefore, if t is a root of $p_{\Gamma}(\cdot)$, the numbers $1, t, \dots, t^{m-1}$, $m = \deg p_{\Gamma}(\cdot)$, are rational independent.

Reducing the components of χ_{Γ} to common denominator, we obtain the following problem: to check that

—for $\Gamma_{(1,1,1,1,1)}$ $q_\Gamma(t) = 1 + t$ is rational independent of 1;
 —for $\Gamma_{(1,1,1,2)}$ $q_\Gamma(t) = (1 + t)(1 + t + t^2)$ is rational independent of $1 + t + t^2$, $1 + t$, $(1 + t)(1 + t)$;
 —for $\Gamma_{(2,2,3)}$ $q_\Gamma(t) = (1 + t + t^2)(1 + t + t^3)$ is rationally independent of $1 + t + t^2 + t^3$, $(1 + t)(1 + t + t^2 + t^3)$, $1 + t + t^2$, $(1 + t)(1 + t + t^2)$, $(1 + t + t^2)(1 + t + t^2)$;
 —for $\Gamma_{(1,3,4)}$ $q_\Gamma(t) = (1 + t)(1 + t + t^2 + t^3)(1 + t + t^2 + t^3 + t^4)$ is rationally independent of $(1 + t + t^2 + t^3)(1 + t + t^2 + t^3 + t^4)$, $(1 + t)(1 + t + t^2 + t^3 + t^4)$, $(1 + t)(1 + t)(1 + t + t^2 + t^3 + t^4)$, $(1 + t + t^2)(1 + t)(1 + t + t^2 + t^3 + t^4)$, $(1 + t)(1 + t + t^2 + t^3 + t^4)$, $(1 + t)(1 + t)(1 + t + t^2 + t^3 + t^4)$, $(1 + t + t^2)(1 + t)(1 + t + t^2 + t^3 + t^4)$, $(1 + t + t^3 + t^4)(1 + t)(1 + t + t^2 + t^3 + t^4)$;
 —for $\Gamma_{(1,2,6)}$ $q_\Gamma(t) = (1 + t)(1 + t + t^2)(1 + t + t^2 + t^3 + t^4 + t^5 + t^6)$ is rationally independent of $(1 + t + t^2)(1 + t + t^2 + t^3 + t^4 + t^5 + t^6)$, $(1 + t)(1 + t + t^2)(1 + t + t^2 + t^3 + t^4 + t^5 + t^6)$, $(1 + t)(1 + t + t^2 + t^3 + t^4 + t^5 + t^6)$, $(1 + t)(1 + t)(1 + t + t^2 + t^3 + t^4 + t^5 + t^6)$, $(1 + t)(1 + t + t^2)$, $(1 + t)(1 + t)(1 + t + t^2)$, $(1 + t + t^2)(1 + t)(1 + t + t^2)$, $(1 + t + t^2 + t^3)(1 + t)(1 + t + t^2)$, $(1 + t + t^2 + t^3 + t^4)(1 + t)(1 + t + t^2)$, $(1 + t + t^2 + t^3 + t^4 + t^5)(1 + t)(1 + t + t^2)$.

But this immediately follows since none of these numbers except contains t^{m-1} , $m = \deg p_\Gamma(\cdot)$. $\square$

The author expresses his deep gratitude to Prof. Yu. S. Samoilenko and Prof. S. V. Popovych for valuable discussions of the subject of this paper.

References

- [1] I. N. Bernstein, I. M. Gelfand, and V. A. Ponomarev, *Coxeter functors and Gabriel's theorem*, Uspekhi Mat. Nauk **28** (1973), no. 2, Russian.
- [2] S. Krugljak, S. Popovych, and Yu. Samoilenko, *Representations of $\ast$ -algebras associated with Dynkin graphs and Horn's problem*, Uchen. Zap. Tavrich. Nat. Univ. **16** (2003), no. 2, 132–139.
- [3] S. A. Krugljak and A. V. Roiter, *Locally scalar representations of graphs in the category of Hilbert spaces*, Funct. Anal. Prilozh. **39** (2005), no. 2, 13–30.
- [4] S. A. Krugljak, V. I. Rabanovich, and Yu. S. Samoilenko, *On sum of projections*, Funct. Anal. Prilozh. **36** (2002), no. 3, 20–35.
- [5] S. A. Kruglyak, *Coxeter functors for a certain class of $\ast$ -quivers and $\ast$ -algebras*, Methods Funct. Anal. Topol. **8** (2002), no. 4, 49–57.
- [6] A. Mellit, *Algebras generated by elements with given spectrum and scalar sum and kleinian singularities*, arXiv:math.RA/0406119.
- [7] A.S. Mellit, *When a sum of three partial reflections is zero*, Ukr. Math. J. **55** (2003), no. 9, 1277–1283.

- [8] A.S. Mellit, V.I. Rabanovich, and Yu.S. Samoilenko, *When a sum of partial reflections is a scalar operator*, Funct. Anal. Prilozh. **38** (2004), no. 2, 91–94.
- [9] V. L. Ostrovskiy, *Representations of algebra associated with Dynkin graph $\tilde{E}_7$* , Ukr. Mathem. J. **56** (2004), no. 9, to appear.
- [10] V. L. Ostrovskiy, *On $*$ -representations of a certain class of algebras related to a graph*, Methods Funct. Anal. Topol. **11** (2005), no. 3, to appear (See also [arXiv:math.RT/0506505](#)).
- [11] I. K. Redchuk, *Separating functions, spectral graph theory and locally scalar representations in Hilbert spaces*, [arXiv:math.RT/0509417](#), 2005.
- [12] I. K. Redchuk and A. V. Roiter, *Singular locally scalar representations of quivers in Hilbert spaces and separating functions*, Ukr. Math. Journ. **56** (2004), no. 6, 947–963.
- [13] Yu. Samoilenko and M. Zavodovsky, *Spectral theorems for $*$ -representations of the algebras $\mathcal{A}_{\Gamma, \chi, com}$ associated with Dynkin graphs*, Methods Funct. Anal. Topol. (2005), no. 1.
- [14] V. S. Shulman, *On representations of limit relations*, Meth. Funct. Anal. Topol. **7** (2002), no. 4, 85–87.
- [15] M. S. Vlasenko, A. S. Mellit, and Yu. S. Samoilenko, *On algebras generated with linearly dependent generators that have given spectra*, Funct. Anal. Appl. **39** (2005), no. 3, to appear.